


Quakes per year. Major = 7-7.9; Great = 8 or larger.

Year	Major quakes	Great quakes	Year	Major quakes	Great quakes
1969	15	1	1989	06	1
1970	20	0	1990	18	0
1971	19	1	1991	16	0
1972	15	0	1992	13	0
1973	13	0	1993	12	0
1974	14	0	1994	11	2
1975	14	1	1995	18	2
1976	15	2	1996	14	1
1977	11	2	1997	16	0
1978	16	1	1998	11	1
1979	13	0	1999	18	0
1980	13	1	2000	14	1
1981	13	0	2001	15	1
1982	10	1	2002	13	0
1983	14	0	2003	14	1
1984	08	0	2004	13	2
1985	13	1	2005	10	1
1986	05	1	2006	9	2
1987	11	0	2007	14	4
1988	08	0	2008	12	0
			2009	16	1
			2010	21	1

<http://earthquake.usgs.gov/earthquakes/eqarchives/year/eqstats.php>

## Gutenberg-Richter Relationship: Magnitude vs. frequency of occurrence



$N(M)$  is number of earthquakes per year (usually) of magnitude  $M$  or LARGER


log is the base 10 log (not ln)

$b$  is usually about 1 for tectonic earthquakes.

If data are for one year, then  $a$  tells us that on average once per year, a quake of magnitude ( $a/b$ ) or bigger happens ( $a$  if  $b = 1$ ).

**How does  $a$  affect the total # of quakes?**


# Gutenberg-Richter Relationship: Magnitude vs. frequency of occurrence



$N(M)$  is number of earthquakes per year (usually) of magnitude  $M$  or LARGER

log is the base 10 log (not ln)

$b$  is about 1 for tectonic earthquakes. It is about 2 for volcanic earthquakes and some earthquake swarms. What does this tell us about the distribution of earthquake sizes on volcanoes?





Here, note that  $M = a/b$  is off to the left (these data are for quakes that happen less than once per year).

What's the magnitude of the once per year quake for each dataset?

Use of these plots: predicting how often big ones occur (we need to know the maximum size)


# Make the G-R plot for worldwide earthquakes





$$\log N(M) = a - bM$$

Southern California Earthquake Center


Why does the curve flatten for small magnitudes?




What governs the high end?




Maximum earthquake size scales with dimension  
of the biggest fault in the area of interest



Moment magnitude versus rupture length





Wells and Coppersmith, 1994



Slip (s) scales with rupture length too, and  $M_0 = GsA$ .  
Therefore  $M_0$  is proportional to  $L^3$  (for small quakes) or  $L^2W$  (bigger ones)


Use fault length and scaling relationships to  
identify biggest quake if faults are mapped

(or we may have records of very large quakes)






Use earthquake statistics and **assumption about maximum earthquake size** if details of faults are not mapped



characteristic earthquake




## characteristic earthquake

on faults with characteristic earthquakes, G-R seismicity statistics work for all but the giant “characteristic earthquake”


this earthquake has a characteristic magnitude and occurs more frequently than GR would suggest

example: Cascadia subduction zone: M9+ earthquakes


Schwartz and Coppersmith (1984)



Youngs and Coppersmith (1985)



People are still arguing about whether the SAF has characteristic earthquakes or not. Seems to depend on which quakes you count (just on the fault? or in some region surrounding the fault, too?) Reason to count off-fault quakes: a big SAF quake could start on another nearby fault (several recent examples!)






# Aftershocks: Omori's Law

$$N(t) = \frac{k}{(t + c)^p}$$

p is approximately 1 (can vary)  
c is small (keeps the denominator above zero)  
k is the number of aftershocks on day one (1st 24 hours)



If k is 100 then 100 on Day 1  
100/2 = 50 on Day 2  
100/3 = 50 on Day 3  
100/4 = 50 on Day 4

What is k on this plot?



How many quakes per day  
one week later, according to  
Omori's Law?

How many quakes per day  
one month later? Consistent  
with the data?


YMMV: different quakes have different aftershock productivity (and sometimes different decay rate)

$$N(t) = \frac{k}{(t + c)^p}$$

p is approximately 1 (can vary)  
c: small number (keeps the denominator above zero)  
k is the number of aftershocks on day one



## Bath's Law: the largest aftershock is 1 magnitude unit smaller than the mainshock



Does not work for every quake but seems to be true on average

Does it work for this one?

Combining GR statistics with Omori's Law gives probability of aftershocks with particular magnitudes, during specific time intervals after a big quake

